

Clinic Well Child Check Guide

Age	Vaccines	Screening	Labs	Other Evaluations
Newborn	Hepatitis B (if not given at birth)			TcB
2-4 weeks		EPDS		Review newborn screen results
2 months	Pediarix (DTaP, IPV, HepB) Hiberix (Hib) Prevnam 15 Rotarix	EPDS		
4 months	Pediarix (DTaP, IPV, HepB) Hiberix (Hib) Prevnam 15 Rotarix	EPDS		
6 months	Pediarix (DTaP, IPV, HepB) Hiberix (Hib) Prevnam 15 *Start flu shots	EPDS		Apply fluoride varnish (first tooth) every 3 months
9 months	Catch up	ASQ-3	Hgb	
12 months	Hep A #1 MMR Varicella		Hgb Lead	
15 months	DTaP Hiberix Prevnam 15			
18 months	Hep A #2 (> 6 months after first dose)	ASQ-3 MCHAT-R		
24 months		ASQ-3 MCHAT-R	Hgb Lead	
30 months		ASQ-3		
3 year		Vision		Start BMI/nutrition/exercise counseling
4 years	Kinrix (DTaP, IPV) Proquad (MMR+Varicella)	Hearing Vision		
5 – 7 years		Hearing Vision		Apply fluoride until 5 year 11 months
8 years		Hearing Vision SCARED-GAD		
9-10 years	9 yo: Gardasil #1	Hearing Vision GAD-SCARED	Lipid 9-11	
11 years	Tdap, Menveo #1 Catch up HPV	GAD-7	Lipid 9-11	
12-14 years		Hearing Vision PHQ-9 CRAFFT GAD-7	Start Hgb (if menstruating female)	HEADSS exam Transition policy
15 years		Hearing Vision PHQ-9 CRAFFT GAD-7	Start offering urine GC/Chlamydia (unless patient declines)	
16 years	Menveo #2 Bexero (2 dose series 6 months apart)	Hearing Vision PHQ-9 CRAFFT GAD-7		HEADSS exam Transition policy
17-21 years		Hearing Vision PHQ-9 CRAFFT GAD-7	Lipid	HEADSS exam Transition policy

- Apply fluoride varnish from eruption of first tooth until 5 years 11 months old UNLESS has received fluoride varnish within the last 3 months
- Start offering flu shots at 6 months old; need 2 doses during first season unless first flu shot is given after age 8 years 11 months old
- Offer Nirsevimab (Beyfortus) at 0-8 months between October-March