

Supplemental Information

SUPPLEMENTAL FIGURE 1. Summary of Down syndrome-specific care.

Action	Pre-natal	Birth up to 1 mo	1 mo up to 1 yr	1 yr up to 5 yr	5 yr up to 12 yr	12 yr up to 21 yr
1. Confirm DS diagnosis with either CVS or amniocentesis prenatally or karyotype postnatally						
2. Review recurrence risk and offer the family referral to a clinical geneticist or genetic counselor.						
3. Offer parent-to-parent and support group information to the family.						
4. Use CDC DS-specific growth charts to monitor weight, length, weight-for-length, head circumference, or BMI. Use standard charts for BMI after age 10 years.		All healthcare visits				
5. Order an echo, to be read by a pediatric cardiologist.						
6. Feeding assessment or video study if any: marked hypotonia, underweight (<5th %ile weight-for-length or BMI), slow feeding or choking with feeds, recurrent or persistent abnormal respiratory symptoms, desaturations with feeds		Any visit				
7. Obtain objective hearing assessment (may be in NBS protocols) and follow EHDJ protocols.			Up to 6 mo			
8. If TM can't be visualized, refer to otolaryngologist for exam with microscope until reliable TM and tympanometry exams are possible		Every 3-6 mo				
9. Car safety seat evaluation before hospital discharge.						
10. CBC with differential		By day 3				
11. If TAM, make caregivers aware of risk/signs of leukemia (e.g., easy bruising/bleeding, recurrent fevers, bone pain)						
12. TSH		At birth (if not in NBS)	Every 5-7 mo	Annually, and every 6 mo if antithyroid antibodies ever detected		
13. RSV prophylaxis based on AAP guidelines.		Annually		Through 2 yr		
14. Discuss cervical spine-positioning for procedures and atlantoaxial stability precautions.		All HMV		Biennially		
15. Assess for CAM use, discourage any unsafe CAM practices.		All HMV				
16. Refer children to early intervention for speech, fine motor or gross motor therapy.		Any visit	Up to 3 yr			
17. If middle ear disease occurs, obtain developmentally-appropriate hearing evaluation.			When ear clear	After treatment		
18. Rescreen hearing with developmentally-appropriate methodology (BAER, behavioral, ear-specific).			Start at 6mo, every 6 mo until established normal bilaterally by ear-specific testing, then annually			
19. Refer to ophthalmologist with experience and expertise in children with disabilities.			By 6 mo			
20. CBC with differential if easy bruising or bleeding, recurrent fevers, or bone pain			Any visit			
21. Assess for sleep-disordered breathing; if present, refer to physician with expertise in pediatric sleep disorders.			At least once by 6 mo, then all subsequent HMV thereafter			
22. Ensure child is receiving developmental therapies, and family understands and is following therapy plan at home.		All HMV				
23. CBC with differential and either (1) a combination of ferritin and CRP, or (2) a combination of serum iron and Total Iron Binding Capacity				Annually		
24. If a child has sleep problems and a ferritin less than 50 mcg/L, the pediatrician may prescribe iron supplement.				Any visit		
25. Vision screening			All HMV, use developmentally-appropriate criteria	Photoscreen (all HMV); if unable, refer to ophthalmologist annually	Photoscreen (all HMV); if unable, refer to ophthalmologist biennially	Visual acuity or photoscreening at all HMV, or ophthalmology-determined schedule
26. If a child has myelopathic symptoms, obtain neutral C-spine plain films (see text for details).				Any visit		
27. Obtain polysomnogram.				Between 3-5 yr		
28. Prepare family for transition from early intervention to preschool.				At 30 mo		
29. Discuss sexual exploitation risks.				At least once	At least once	At least once
30. Make developmentally-appropriate plans for menarche, contraception (advocate/offer LARC), and STI prevention.					As developmentally-appropriate, then all subsequent HMV	
31. Discuss risk of DS if patient were to become pregnant.					At least once	At least once
32. Assess for any developmental regression.			All HMV			
33. Discuss and facilitate transitions: education, work, finance, guardianship, medical care, independent living					All HMV starting at 10 yr	

	Do once at this age	Abbreviations: DS, Down syndrome; CVS, Chorionic villus sampling; HMV, Health Maintenance Visit; BMI, Body mass index; CDC, Centers for Disease Control; EHDJ, Early Hearing Detection and Intervention; NBS, Newborn screen; CAM, Complementary and alternative medicine; BAER, Brainstem auditory evoked response; TM, Tympanic membrane; TAM: transient abnormal myelopoiesis
	Do if not done previously	
	Repeat at indicated intervals	
	See report for end point	